



Announcement

Guidance for Relief Workers and Others Traveling to Haiti for Earthquake Response

This information is current as of today, January 28, 2010 at 08:16 EST

Updated: January 27, 2010

This notice is to advise relief workers and other personnel traveling to Haiti to assist with the humanitarian response following the January 12th earthquake near Port-au-Prince. Conditions in the area remain hazardous, including extensive damage to buildings, roads, and other infrastructure.

Before You Depart for Haiti

Recommended Vaccines

A number of vaccines are recommended for travelers to Haiti. See your doctor before you travel to make sure you have had all necessary vaccines.

- **Routine** (<http://www.cdc.gov/vaccines/recs/schedules/default.htm>): Be sure that you are up to date on vaccines such as measles/mumps/rubella (MMR), diphtheria/pertussis/tetanus (DPT), polio, seasonal and H1N1 flu, and varicella. It is especially important to have a current tetanus shot.
- **Hepatitis A or immune globulin (IG)** (</travel/yellowbook/2010/chapter-2/hepatitis-a.aspx>): Even if your departure is imminent, one dose of hepatitis A vaccine provides adequate short-term protection for healthy people. For long term protection, a second dose is required 6–18 months after the first dose, depending on the brand of vaccine used.
- **Typhoid** (</travel/yellowbook/2010/chapter-2/typhoid-paratyphoid-fever.aspx>): There are 2 vaccines available for typhoid prevention. The injectable vaccine may be preferable to the oral vaccine in cases where travel is imminent. The oral vaccine requires refrigeration and 4 tablets taken every other day over one week.
- **Hepatitis B** (</travel/yellowbook/2010/chapter-2/hepatitis-b.aspx>): If your departure is imminent, the first in a 3-dose series (day 0, 1 month and 6 months) may provide some protection. An accelerated dosing schedule may be used (doses at days 0, 7, and at 21–30 days with a booster at 12 months).
- **Rabies** (</travel/yellowbook/2010/chapter-2/rabies.aspx>): If your activities in Haiti will bring you into contact with animals such as dogs, cats, bats, raccoons, or skunks, you should consider pre-exposure rabies vaccination, which is a 3-shot series (days 0, 7, and 21 or 28) given before travel. Even if you receive pre-exposure vaccination, you will still need immediate medical treatment if you are bitten or scratched by an animal. (See the [Animals \(#animals\)](#) section for more information.)

Insect-borne Diseases

Malaria

Malaria (<http://www.cdc.gov/malaria/index.htm>) occurs in all parts of Haiti. Ways to prevent malaria include the following:

- Taking a prescription antimalarial drug
- Using insect repellent and wearing long pants and sleeves to prevent mosquito bites
- Sleeping in air-conditioned or well-screened rooms or using bed nets

No antimalarial drug is 100% protective, so it is important to use all three ways to prevent malaria.

All of the following antimalarial drugs are equal options for preventing malaria in Haiti: Atovaquone/proguanil (Malarone), chloroquine, doxycycline, or mefloquine. Each drug has its own side effects, contraindications, and precautions. You will need to talk to your doctor to decide which of these drugs would be best for you, depending on your current health, medical history, drug allergies, and specific needs. Additional information can be found on the [Drugs to Prevent Malaria \(http://www.cdc.gov/malaria/control_prevention/drugs.htm\)](http://www.cdc.gov/malaria/control_prevention/drugs.htm) page.

Malaria is always a serious disease and may be a deadly illness. If you become ill with a fever or flu-like illness either while in Haiti or after you return home (for up to 1 year), you should seek immediate medical attention and should tell the doctor that you have recently been in Haiti.

Dengue

Dengue is a common infection in Haiti. Dengue is a disease caused by a virus transmitted to people by the bite of an infected mosquito. Some important information to know about dengue as you travel to Haiti:

- No vaccine or medications are available to prevent dengue.
- The best way to reduce your risk of dengue is to protect yourself from mosquito bites (see the section below called “[Protection Against Insects and Animals \(#insects\)](#)”).
- The mosquitoes that spread dengue usually bite at dusk and dawn but may bite at any time during the day.
- Symptoms and signs are high fever, chills, headache and muscle pain. Additionally, a faint rash on the trunk and upper arms may appear on the second to third day of illness.
- There are no specific medicines to treat dengue, so treatment is supportive with fever-reducing medicines and fluids.
- You can help control mosquito populations by draining all standing water that you find in open containers left outdoors.
- If you are in the Dominican Republic awaiting entry into Haiti, be aware that dengue is also common there.

Other Infectious Diseases

HIV: Haiti has a high prevalence of HIV infection. To reduce the risk of HIV and other sexually transmitted diseases, always use latex condoms. Healthcare workers should also take the following additional precautions:

- Wear gloves for touching blood and body fluids, mucous membranes, or broken skin and for handling items or surfaces soiled with blood or body fluids.
- Use masks and protective eyewear or face shields to prevent exposure of the mouth, nose, and eyes during procedures that are likely to generate droplets of blood or body fluids.
- Wear gowns or aprons during procedures that are likely to generate splashes of blood or body fluids.

Tuberculosis (TB): Rates of tuberculosis are very high in Haiti.

- If you anticipate giving medical care to or working closely with ill or injured victims or other displaced persons in Haiti, a [tuberculin skin test](http://www.cdc.gov/tb/publications/factsheets/testing/skintesting.htm) (<http://www.cdc.gov/tb/publications/factsheets/testing/skintesting.htm>) or TST (ideally, a two-step test) is recommended before travel and then 8–10 weeks after return.
 - If you have had a negative result on a tuberculin skin test within the past 12 months, you do not need to be tested before going to Haiti.
 - If you have had a positive result on a tuberculin skin test in the past, you do not need to be tested again.
- If you are in contact with known TB patients or persons suspected of having TB, use a personal respiratory protective device (e.g., N-95 [respirator](http://www.cdc.gov/niosh/npptl/topics/respirators/disp_part/RespSource3.html) (http://www.cdc.gov/niosh/npptl/topics/respirators/disp_part/RespSource3.html)).
- **Regardless of tuberculin skin test results, any person who develops [symptoms of TB](http://www.cdc.gov/tb/topic/basics/signsandsymptoms.htm) (<http://www.cdc.gov/tb/topic/basics/signsandsymptoms.htm>) during or after deployment should see a doctor immediately.** Symptoms of TB can include a bad cough that lasts 3 weeks or longer; pain in the chest; coughing up blood or sputum (phlegm from deep inside the lungs); weakness or fatigue; weight loss; no appetite; chills; fever; and sweating at night.

Anthrax: Anthrax occurs in Haiti and is primarily transmitted by direct contact with infected animals or with contaminated products from infected animals. Cases of cutaneous and inhalation anthrax have been reported among the local population.

- Cutaneous (skin) anthrax infection usually begins as a small sore that develops into a blister. The blister then develops into a skin ulcer with a black area in the center. The sore, blister, and ulcer do not hurt.
- The first symptoms of inhalation anthrax are like cold or flu symptoms and can include a sore throat, mild fever (>100° F), and muscle aches. Later symptoms include cough, chest discomfort, shortness of breath, tiredness, and muscle aches.

If you develop any of these symptoms, see a healthcare provider immediately.

Key Items to Bring

There will be almost no infrastructure support available in Haiti for the immediate future. Relief workers, volunteers, and other travelers will need to be self-sufficient.

Pack basic supplies, including

- Sleeping in beds covered by a bed net (preferably treated with permethrin), if not sleeping in an air-conditioned or well-screened room.
- Spraying rooms with products effective against flying insects, such as those containing pyrethroid.
- For detailed information about insect repellent use, see [Insect and Arthropod Protection \(/travel/content/insectprotection.aspx\)](/travel/content/insectprotection.aspx).

Animals

Direct contact with animals can spread diseases like rabies or cause serious injury or illness. Displaced animals may revert to the wild and go about in packs. They will also be hungry and may be searching for food and may be more likely to bite. Because you may not have access to appropriate medical care, it is important to prevent animal bites and scratches.

- Stay away from all animals, including dogs and cats. Even animals that look like healthy pets can have rabies or other diseases.
- If you are bitten or scratched, wash the wound well with soap and clean water and **seek medical care right away**. If you have a povidone-iodine solution (such as Betadine), use that to clean the wound after washing it.
 - If you have **not** received pre-exposure rabies vaccination: You will need shots of rabies immune globulin (or RIG) and 4 shots of rabies vaccine over 1 month.
 - If you have received pre-exposure rabies vaccination: You will need two shots of rabies vaccine over 3 days.
 - Rabies vaccine and rabies immune globulin are probably not available in Haiti at this time. You will need to have a plan in place to fly to another country for treatment.
- Resist the urge to rescue animals with the intent to bring them home to the United States. Dogs and cats may be infected with rabies but not show signs until several days or weeks after you first encounter them.
- After you return from Haiti, be sure to tell your doctor or state health department if you were bitten or scratched during travel.
- For more information about how to protect yourself from other risks related to animals, see [Animal-Associated Hazards \(/travel/yellowbook/2010/chapter-2/animal-associated-hazards.aspx\)](/travel/yellowbook/2010/chapter-2/animal-associated-hazards.aspx). To learn more about Rabies see [CDC's Rabies homepage \(http://www.cdc.gov/rabies\)](http://www.cdc.gov/rabies).

Injury

The risk of injury after an earthquake is high. Hazards such as electrocution from downed power lines and structural damage to buildings and roads all pose a risk. Accidents and violence are documented risks for humanitarian workers and cause more deaths than disease and natural causes. According to the World Health Organization, injuries are among the leading causes of preventable death in travelers.

- The majority of the road network in Haiti is not paved. Haiti is predominately mountainous and has extensive deforestation and soil erosion, making travel over roadways especially hazardous. Exercise extreme care when traveling on roads particularly in rural areas.
- There has been extensive structural damage to buildings in Haiti. Avoid unstable structures if possible.
- Other potential hazards to be aware of in collapsed buildings include standing water from water system breaks, natural gas leaks, airborne smoke and dust, hazardous materials such as ammonia or leaking fuels, exposure to germs from sewer line breaks, and exposed wiring.
- Use personal protection equipment, such as hard hats and steel-toed boots, if in areas with damaged buildings.

Exposure to Human Remains (<http://emergency.cdc.gov/disasters/handleremains.asp>)

Human remains may contain blood-borne viruses and diarrhea-causing bacteria. Relief workers who are handling remains should take precautions to avoid being exposed to these organisms.

- Protect your face from splashes of body fluids and fecal material by using a plastic face shield or a combination of eye protection and surgical mask. In extreme situations, a cloth tied over the nose and mouth can be used to block splashes.
- Protect your hands from direct contact with body fluids and from injuries that break the skin by using a combination of a cut-proof inner layer glove and a latex (or similar) outer layer.
- Wash your hands with soap and water or with an alcohol-based hand cleaner immediately after you remove the gloves.
- Protect your feet and ankles against sharp debris by wearing foot wear that covers the entire foot and has thick soles.
- Give prompt care—including immediate cleansing with soap and water, and a tetanus booster if indicated—to anyone who is injured during work with human remains.

For more extensive information about working with human remains after a disaster, see the [Interim Health Recommendations for Workers who Handle Human Remains After a Disaster \(http://emergency.cdc.gov/disasters/handleremains.asp\)](http://emergency.cdc.gov/disasters/handleremains.asp) fact sheet.

Psychological/Emotional Difficulties

As a first responder or relief worker, you may encounter extremely stressful situations, such as witnessing a tremendous loss of life, serious injuries, missing and separated families, and destruction of whole areas. It is important to recognize that these experiences may cause you psychological or emotional difficulties.

Some Common Normal Reactions to a Disaster

- Profound sadness, grief, and anger.
- Not wanting to leave the scene until the work is finished.
- Trying to override stress and fatigue with dedication and commitment.
- Denying the need for rest and recovery time.

Ways to Help Manage Your Stress

- Limit on-duty work time to no more than 12 hours per day.
- Rotate work assignments between high stress and lower stress functions.
- Drink plenty of water and eat healthy snacks and energy foods.
- Take frequent, brief breaks from the scene when you are able.
- Keep an object of comfort with you such as a family photo, favorite music, or religious material.
- Stay in touch with family and friends.
- Pair up with another responder so that you can monitor one another's stress.

To learn about mental health resources, see:

- [Self-Care Tips for Stress \(http://mentalhealth.samhsa.gov/publications/allpubs/KEN-01-0098/\)](http://mentalhealth.samhsa.gov/publications/allpubs/KEN-01-0098/)
- Download Podcasts on stress management for first responders: [Stress Management for Emergency Responders - What Responders Can Do \(http://www2c.cdc.gov/podcasts/player.asp?f=10613\)](http://www2c.cdc.gov/podcasts/player.asp?f=10613)
- [NIOSH: Traumatic Incident Stress: Information for Emergency Response Workers \(http://www.cdc.gov/niosh/docs/2002-107/\)](http://www.cdc.gov/niosh/docs/2002-107/)
- [NIOSH: Estrés por sucesos traumáticos: Información para el personal de Emergencia \(http://www.cdc.gov/spanish/niosh/docs/unp-trinstrsSP.html\)](http://www.cdc.gov/spanish/niosh/docs/unp-trinstrsSP.html)

After You Come Home

If you are not feeling well, you should see your doctor and mention that you have recently returned from response and relief work in Haiti. Also tell your doctor if you were bitten or scratched by an animal while traveling.

Symptoms of malaria can develop up to one year after travel, so be alert for fever or flu-like symptoms.

Approximately one-third of aid workers report depression shortly after returning home, and more than half of returned aid workers have reported feeling predominantly negative emotions on returning home, even though many reported that their time overseas was positive and fulfilling. You might want to see a mental health professional to help you adjust back into your home environment.

More Information

- [Haiti destination page \(/travel/destinations/haiti.aspx\)](/travel/destinations/haiti.aspx) on the Travelers' Health website
- [Haiti country specific information \(/travel/forward.aspx?faHR0cDovL3d3dy50cmF2ZWwuc3RhZGUuZ292L3RyYXZlbc9jaXNfcGFfdHcvY2l2L2Npc18xMTM0Lmh0bWw%3d-KHXgbb71bjI%3d\)](/travel/forward.aspx?faHR0cDovL3d3dy50cmF2ZWwuc3RhZGUuZ292L3RyYXZlbc9jaXNfcGFfdHcvY2l2L2Npc18xMTM0Lmh0bWw%3d-KHXgbb71bjI%3d) from the US Department of State
- [United Nations Relief Web \(/travel/forward.aspx?faHR0cDovL3d3dy5yZWxpZWZ3ZWluaW50L3JlL2RiYy5uc2YyZG9jMTA4P09wZW5Gb3JtJmFtcDtlbWlkPUVRLTIwMTAtMDAwMDA%3d%3d-zPZ6oCqeA6U%3d\)](/travel/forward.aspx?faHR0cDovL3d3dy5yZWxpZWZ3ZWluaW50L3JlL2RiYy5uc2YyZG9jMTA4P09wZW5Gb3JtJmFtcDtlbWlkPUVRLTIwMTAtMDAwMDA%3d%3d-zPZ6oCqeA6U%3d)

- Page last reviewed: January 16, 2010
- Page last updated: January 27, 2010
- Page created: January 14, 2010
- Content source:
Division of Global Migration and Quarantine
National Center for Preparedness, Detection, and Control of Infectious Diseases



Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, 24 Hours/Every Day - cdcinfo@cdc.gov